

ANNEX 1

INTEGRATED ADVICE – 25-29 AND 30-64 YEAR OLDS

Age range: 25-29 years					
Result cytological examination	Result reflex hrHPV test	(Integrated) advice	Result 2 nd triage (in 12 months)	Result reflex hrHPV test	(integrated) advice
NILM	NA	Regular screening interval (in 3 calendar years)			
ASC-US	--> hrHPV negative	Result of reflex HPV test, with recommendation, will follow Regular screening interval (in 3 calendar years)			
	--> hrHPV positive	Repeat cytology in 12 months	--> NILM	NA	Regular screening interval (in 3 calendar years)
			--> ASC-US		Result of reflex HPV test, with recommendation, will follow Regular screening interval (in 3 calendar years)
				--> hrHPV negative	Regular screening interval (in 3 calendar years)
				--> hrHPV positive	Immediate referral for colposcopic examination
			--> ≥ LSIL		Immediate referral for colposcopic examination
LSIL	NA	Repeat cytology in 12 months	--> NILM	NA	Regular screening interval (in 3 calendar years)
			--> ASC-US		Result of reflex HPV test, with recommendation, will follow Regular screening interval (in 3 calendar years)
				--> hrHPV negative	Regular screening interval (in 3 calendar years)
				--> hrHPV positive	Immediate referral for colposcopic examination
			--> ≥ LSIL		Immediate referral for colposcopic examination
≥ ASC-H/AGC	NA	Immediate referral for colposcopic examination		NA	
INSU	NA	New sampling after 6 weeks at the earliest		NA	
Age range: 30-64 years					
Result hrHPV test	Result reflex cytology	(Integrated) advice	Result 2 nd triage (in 12 months)	(Result reflex cytology)	Advice
hrHPV negative	NA	Regular screening interval (in 5 calendar years)			
hrHPV non-16/18 positive		Result of reflex cytology, with recommendation, will follow Immediate referral for colposcopic examination			
	--> ≥ ASC-US	Repeat hrHPV testing in 12 months	--> hrHPV negative		Regular screening interval (in 5 calendar years)
	--> NILM		--> hrHPV positive	(cytology, not as triage)	Immediate referral for colposcopic examination
					Result of reflex cytology will follow
HPV 16/18 positive	(cytology, not as triage)	Immediate referral for colposcopic examination			
		Result of reflex cytology will follow			
HPVi		New sampling after 6 weeks at the earliest			

≥ASC-H/AGC = ASC-H or a more severe abnormality (HSIL, SCC, AIS, AC) or AGC

INSU: insufficient cytology, not representative sample (lack of epithelial cells/insufficient cellular material, cell lysis, abundant blood, ..)

HPVi: inconclusive HPV test result

NB: Immediate referral for colposcopic examination is understood within 3 months or faster, according to the severity of the screen-positive result (cfr. 5.3.1)

Cave: If on cytological examination **normal** endometrial cells are found in an entitled person > 45 years, additional advice is given: 'Correlation with clinic is indicated to exclude endometrial pathology in post-menopausal women'.

Cave: If on cytological examination **abnormal** endometrial cells are found at any age, additional advice is given: 'Exploration to exclude endometrial pathology'.

In bold: the actual integrated advice, after samples were send for reflex testing.